



Fates of methane from different lake habitats: Connecting whole-lake budgets and CH₄ emissions

David Bastviken,¹ Jonathan J. Cole,² Michael L. Pace,² and Matthew C. Van de Bogert³

Received 27 September 2007; revised 3 February 2008; accepted 28 February 2008; published 24 May 2008.

[1] Methane (CH₄) represents a major product of organic matter decomposition in lakes. Once produced in the sediments, CH₄ can be either oxidized or emitted as a greenhouse gas to the atmosphere. Lakes represent an important source of atmospheric CH₄, but the relative magnitudes of the internal pathways that lead to CH₄ emissions are not yet clear. We quantified internal cycling and methane emissions in three lakes during summer stratification. These methane budgets included: sediment release of CH₄ at different depths; water column transport patterns and methane oxidation; methane storage in the water column; and methane emissions to the atmosphere by diffusion and ebullition. The contribution of CH₄ carbon, via oxidation by methanotrophic bacteria, to pelagic food webs was also estimated. Despite the very low concentration of CH₄ in surface waters, shallow, epilimnetic sediments were major contributors of CH₄ to the atmosphere. While 51–80% of the CH₄ produced in deep sediments was oxidized in the water column, most of the CH₄ released from shallow sediment escaped oxidation and reached the atmosphere. Epilimnetic sediments accounted for 100% of CH₄ emitted during summer stratification, and 14–76% considering the release of CH₄ stored in deep water layers during lake circulation after the stratification period; diffusive emission accounted for 26–48% and ebullition the remainder. These results indicate that it is important to address transport rates of CH₄ from the shallow sediment along with the production-consumption processes when trying to understand methane dynamics and the regulation of lake methane emissions.

Citation: Bastviken, D., J. J. Cole, M. L. Pace, and M. C. Van de Bogert (2008), Fates of methane from different lake habitats: Connecting whole-lake budgets and CH₄ emissions, *J. Geophys. Res.*, *113*, G02024, doi:10.1029/2007JG000608.